



## Consideration of the Webb-Pearman-Leuning theory for closed-path eddy covariance measurements: What should be done?

Ibrom, Andreas; Massman, W.J.

*Published in:*  
Geophysical Research Abstracts

*Publication date:*  
2008

*Document Version*  
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

*Citation (APA):*  
Ibrom, A., & Massman, W. J. (2008). Consideration of the Webb-Pearman-Leuning theory for closed-path eddy covariance measurements: What should be done? *Geophysical Research Abstracts*, 10, EGU2008-A-04560.

---

### General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.



## **Consideration of the Webb-Pearman-Leuning theory for closed-path eddy covariance measurements: What should be done?**

**A. Ibrom** (1), W. J. Massman (2)

(1) Biosystems Department, Risø National Laboratory, Denmark, Technical University of Denmark (DTU), 4000 Roskilde Denmark, [andreas.ibrom@risoe.dk](mailto:andreas.ibrom@risoe.dk) / Fax: ++45 46773160 / Phone: ++45 46774177

2) USDA/Forest Service, Rocky Mountain Research Station, Fort Collins, CO 80526, USA, [wmassman@fs.fed.us](mailto:wmassman@fs.fed.us) / Fax: +1 970-4981314 / Phone: ++1 970-4981296

When calculating turbulent fluxes from scalar densities, density fluctuations due to sensible heat and water vapour fluxes have to be considered (Webb et al. 1980, WPL). The original work only considered open-path eddy covariance (EC) systems - we discuss here the effects for closed-path EC systems.

Contrary to open-path sensors, which measure scalar density more or less unchanged, a closed-path EC system acts as a physical low-pass filter to the atmospheric scalar concentration signal. Large part of the low-pass filtering effect is caused by gas specific, microphysical or/and chemical reactions of the scalar with the surfaces of the EC system. This is why cut-off frequencies differ for different atmospheric scalars.

Whereas temperature fluctuations are effectively dampened in the system, fluctuations of water vapour still contaminate scalar measurements ('dilution effect'). The problem arises how to correct for the dilution effect given the scalar specific degree of low-pass filtering in a closed-path EC system. During the last years, four different recommendations were given with regard to the procedure how to correct for the dilution effect:

1<sup>st</sup> Applying the spectral corrections to all relevant measured scalar concentrations prior to applying the WPL theory (as, e.g. mentioned in an unpublished erratum to

Leuning (2007),

2<sup>nd</sup> correcting only for passive spectral attenuation as opposed to active attenuation by the EC system before applying the WPL theory (Massman 2004),

3<sup>rd</sup> avoiding the necessity for correction of the dilution effect by converting mole fractions ( $\chi$ ) into molar mixing ratios ( $r$ ) at the raw data level (Ibrom et al. 2007) or,

4<sup>th</sup>, using the uncorrected covariance of  $\chi_{H_2O}$  with the vertical wind speed that is calculated at the same time lag as for the scalar in consideration in the WPL correction (Ibrom et al. 2007).

At the example of a data set of closed-path EC CO<sub>2</sub> flux measurements from the Sorø beech forest EUROFLUX site, we investigate the uncertainties and inaccuracies that are introduced by either way of consideration. From this, recommendations will be derived, how to best consider dilution effects for CO<sub>2</sub> flux measurements in closed-path EC systems.

#### References:

Ibrom, A., E. Dellwik, S.E. Larsen and K. Pilegaard. 2007. On the use of the Webb - Pearman - Leuning - theory for closed-path eddy correlation measurements. *Tellus B*. 59B:937-946.

Leuning, R. 2007. The correct form of the Webb, Pearman and Leuning equation for eddy fluxes of trace gases in steady and non-steady state, horizontally homogeneous flows. *Boundary-Layer Meteorol.* 123:263-267.

Massman, W.J. 2004. Concerning the measurement of atmospheric trace gas fluxes with open- and closed-path eddy covariance system: The WPL terms and spectral attenuation. *In Handbook of Micrometeorology* Eds. X. Lee, W.J. Massman and B. Law. Kluwer Academic Publishers, Dordrecht, Boston, London, pp. 133-160.

Webb, E.K., G.I. Pearman and R. Leuning. 1980. Correction of flux measurements for density effects due to heat and water vapour transfer. *Quart J Roy Meteorol Soc.* 106:85-100.